



3-10-2008

Anisotropic Light Scattering from Ferrofluids

Corneliu Rablau

Prem Vaishnava

Ratna Naik

Gavin Lawes

Ronald Tackett

Follow this and additional works at: https://digitalcommons.kettering.edu/physics_conference



Part of the [Physics Commons](#)

NASA/ADS

Anisotropic Light Scattering from Ferrofluids

Rablau, Corneliu; Vaishnava, Prem; Naik, Ratna; Lawes, Gavin; Tackett, Ron; Sudakar, C.

We have investigated the light scattering in DC magnetic fields from aqueous suspensions of Fe_3O_4 nanoparticles coated with tetra methyl ammonium hydroxide and $\gamma\text{-Fe}_2\text{O}_3$ nanoparticles embedded in alginate hydrogel. For Fe_3O_4 ferrofluid, anomalous light scattering behavior was observed when light propagated both parallel and perpendicular to the magnetic fields. This behavior is attributed to the alignment and aggregation of the nanoparticles in chain-like structures. A very different light scattering behavior was observed for $\gamma\text{-Fe}_2\text{O}_3$ alginate sample where, under the similar conditions, the application of the magnetic field produced no structured change in scattering. We attribute this difference to the absence of chain-like structures and constrained mobility of iron nanoparticles in the alginate sample. The observation is in agreement with our relaxation and dissipative heating results^{^1} where both samples exhibited Neel relaxation but only the Fe_3O_4 ferrofluid showed Brownian relaxation. The results suggest that Brownian relaxation and nanoparticle mobility are important for producing non-linear light scattering in such systems. ^{^1}P.P. Vaishnava, R. Tackett, A. Dixit, C. Sudakar, R. Naik, and G. Lawes, J. Appl. Phys. 102, 063914 (2007).

Publication:


American Physical Society, 2008 APS March Meeting, March 10-14, 2008, abstract id. P24.013

Pub Date:

March 2008

Bibcode:

2008APS..MARP24013R

 Feedback/Corrections? (http://adsabs.harvard.edu/adsfeedback/submit_abstract.php?bibcode=2008APS..MARP24013R)